



8401 West Dodge Road • Suite 115 • Omaha, NE 68114 • 1-800-222-1222 • 402-955-5555 • www.NebraskaPoison.com

Serving Nebraska, Idaho, American Samoa, and the Federated States of Micronesia

From: Ron Kirschner, MD, Medical Director
To: ALL HEALTH CARE PROFESSIONALS
Subject: Emergency department naloxone dosing and observation time after administration
Date: 2/28/17

- Naloxone (NAL) is a potentially life-saving opioid antagonist that can reverse respiratory depression and obviate the need for intubation.
- The starting dose recommended in common medical references ranges from < 0.05 mg IV to more than ten-fold greater (Connors).
- A small retrospective study (n=15) found that the median dose needed to reverse respiratory compromise in emergency department (ED) patients was 0.08 mg (range 0.04-0.12) (Kim).
- Higher doses can precipitate opioid withdrawal leading to multiple complications. For patients with respiratory compromise and possible opioid dependence, we recommend starting with 0.04 mg IV.
- In small children, especially those with buprenorphine exposure, higher NAL dosing (e.g. 0.1 mg/kg IV) is appropriate.
- Because NAL's duration of action is short (30-90 minutes) CNS/respiratory depression may recur. The appropriate observation time following EMS or ED administration has been a subject of debate.
- Previous studies of NAL reversal in parenteral heroin abusers suggested that recurrence was unlikely if alert and without respiratory compromise after one hour (Willman).
- Because heroin substitution with fentanyl analogs is increasingly common (Gladden), and we have less NAL reversal experience in this situation, we recommend at least 4 hours of observation following NAL treatment for parenteral opioid abuse.
- Patients with NAL reversal following extended release opioid or methadone ingestion require longer observation.
- For patients with recurrent respiratory depression, continuous naloxone infusion can be initiated at 2/3 the effective reversal dose per hour and titrated to effect (Goldfrank).

References

Connors NJ. The evolution of recommended naloxone dosing for opioid overdose by medical specialty. *J Med Toxicol* 2016; 12: 276.
Gladden R. Fentanyl law enforcement submissions and increases in synthetic opioid-involved overdose. *MMWR* 2016 Aug 26; 65: 644.
Goldfrank L. A dosing nomogram for continuous infusion intravenous naloxone. *Ann Emerg Med* 1986; 15: 566.
Kim HK. Reversal of opioid-induced ventilatory depression using low dose naloxone (0.04 mg). *J Med Toxicol* 2016; 12: 107.
Willman MW. Do heroin overdose patients require observation after receiving naloxone? *Clin Toxicol* 2017; 55: 81.

**Our certified nurse specialists in poison information and physician toxicologists
are available 24 hours a day to answer your questions.**