



From: Ron Kirschner, MD, Medical Director

To: ALL HEALTH CARE PROFESSIONALS

Subject: Emergency department naloxone dosing and observation time after administration

Date: 01/13/22

- Naloxone (NAL) is a potentially life-saving opioid antagonist that can reverse respiratory depression and obviate the need for intubation.
- The starting dose recommended in common medical references ranges from < 0.05 mg IV to more than ten-fold greater (Connors).
- A small retrospective study (n=15) found that the median dose needed to reverse respiratory compromise in emergency department (ED) patients was 0.08 mg (range 0.04-0.12) (Kim).
- Higher doses can precipitate opioid withdrawal leading to multiple complications. For patients with respiratory compromise and possible opioid dependence, we recommend starting with 0.04 mg IV.
- In small children, especially those with buprenorphine exposure, higher NAL dosing (e.g. 0.1 mg/kg IV) is appropriate.
- Because NAL's duration of action is short (30-90 minutes) CNS/respiratory depression may recur. The appropriate observation time following EMS or ED administration has been a subject of debate.
- Patients with NAL reversal following extended release opioid or methadone ingestion require longer observation.
- For patients with recurrent respiratory depression, continuous naloxone infusion can be initiated at 2/3 the effective reversal dose per hour and titrated to effect (Goldfrank).

Call the Poison Center 24 hours a day. Dial 288, wait for operator, then dial 888-222-4516. You will speak directly with a Registered Nurse or Pharmacist 24/7/465.